

## New taxa in genus *Dactylorhiza* NECK. ex NEVSKI and new localities in family *Orchidaceae* to Hungary

VLČKO, Jaroslav<sup>1</sup> – HRIVNÁK, Richard<sup>1</sup> – BALÁZS Pavol<sup>2</sup>

(1) Department of Phytology, Faculty of Forestry, Technical University in Zvolen, T.G. Masaryka 24, SK-960 53 Zvolen, Slovakia

(2) Department of Ecology, Faculty of Academic and Natural Sciences, SK-081 16 Prešov, Slovakia

The group *Dactylorhiza incarnata* is, in the European context, rich in species, and taxonomically not completely clarified in some aspects. The following taxa are known to occur at the territory of Central Europe: *Dactylorhiza incarnata* subsp. *incarnata* L., *D. incarnata* subsp. *haematodes* (Rchb.) Soó, *D. pulchella* (Druce) Averyanov, *D. cruenta* and *D. ochroleuca*. In Hungary, the taxa *Dactylorhiza incarnata* subsp. *incarnata* and *D. ochroleuca* have been recorded until now (BORSOS 1960, Soó 1973). The same taxa are given in MOLNÁR et al. (1995). Despite the decreased number of localities, where *Dactylorhiza incarnata* subsp. *incarnata* occurs, large populations of this species still exist.

During a three-week stay in NE-Hungary in June 1996, which was organized on the basis of an agreement between the Slovak Environmental Agency and the Directorate of the Bükk National Park, we visited several localities where interesting species of the *Orchidaceae* family occur. Montane meadows at the village Regéc in the Zempléni Hegység Mts. belonged to the most interesting ones.

The this locality, we recorded the following representatives of the *Orchidaceae* family: *Platanthera bifolia* subsp. *latiflora*, *Gymnadenia conopsea* subsp. *conopsea*, *Traunsteinera globosa*, *Dactylorhiza fuchsii* subsp. *soóana*, *D. majalis* subsp. *majalis*, *Listera ovata*, *D. sambucina* (confirming SIMON 1977a, p. 39., SIMON 1977b, MATUS 1997) and *D. incarnata* subsp. *incarnata* which is new to the territory.

The impulse for presenting this study was the fact that we succeeded in finding two taxa which are new for the Hungarian flora, namely *Dactylorhiza incarnata* subsp. *haematodes* and *D. pulchella*, as well as rarely occurring hybrid *D. × aschersoniana*.

### Methods

The scientific names of higher plants follow MARHOLD & HINDÁK (1998), names of plant communities follow ELLMAUER & MUCINA (1993). *Dactylorhiza pulchella* was classified following DELFORGE (1994), *D. × aschersoniana* nsubsp. *wiśniewski* following POTŮČEK (1990). For phytosociological records, scale of abundance and dominance (BARKMAN et al. 1964) was used, where the value 2m represents a high abundance and a dominance of 5 %, value 2a represents dominance of 5 to 12,5 % and 2b 12,5 to 25 %.

### *Dactylorhiza incarnata* subsp. *haematodes* (Rchb.) Soó

*D. incarnata* subsp. *haematodes* differs from the nominate subspecies by brown-black maculate leaves, and frequently by flowers of a darker colour. The leaves can be spotted on one side in case of var. *haematodes* (Reichenb.) Soó, or on both sides in case of var. *hyphaematodes* (Neuman) Landw. The spots are mostly concentrated in the upper part of leaves. On the front side, the spots are substantially larger, but less numerous than on the back side. In some populations, specimens occur which are maculate on both sides of leaves almost continuously. At the beginning of flowering period, spots are dark, but they rapidly become pale.

In Central Europe, this taxon can be erroneously classified as *Dactylorhiza cruenta*. That species is characterized by smaller height, thicker stalk, less leaves (3 to 5), which are shorter, wider, maculate on both sides, more declining from the stalk, and obtusely grooved lengthwise. The inflorescence is shorter, with less flowers. The painting at the labellum is more narrow and articulated.

On the Regéc locality, we found three individuals with leaves spotted on both sides, identified as *Dactylorhiza incarnata* subsp. *haematodes* var. *hyphaematodes*.

In Germany and Slovakia, also the hybrid with *Dactylorhiza majalis* was found, namely *D. × aschersoniana* nsubsp. *wiśniewski* (Hemke) Potůček (POTŮČEK 1990). In contrast to *D. × aschersoniana*, it is spotted also on the back side of leaves, but the spots are mostly less numerous and small. Due to this fact, it can easily be neglected.

***Dactylorhiza pulchella* (Druce) Averyanov**(Syn. *D. incarnata* subsp. *serotina* (Hausskn.) D. M. Moore et Soó)

As compared to *Dactylorhiza incarnata* subsp. *incarnata*, it is characterized by smaller height (10 to 25 cm), rarely up to 50 cm, smaller number (3 to 5) of more slender leaves, few flowers in the inflorescence, and at lower altitudes, flowering time is postponed by one or two weeks.

At the locality near the village Regéc, we found one individual, near the village Nyirád, we found together with A. MOLNÁR V. and R. VIDÉKI cca. five individuals in 1998.

*Dactylorhiza pulchella* occurs frequently together with *D. incarnata* subsp. *incarnata*, or with the other taxa of the genus *Dactylorhiza*. An extensive hybridization, even backward hybridization is frequent, what complicates the determination of individual plants.

***Dactylorhiza* × *aschersoniana* (Hausskn.) Borsos et Soó**

This is a hybrid between *D. incarnata* and *D. majalis*, and it is very variable. Intermediate individuals are simply to be determined. In case of a hybrid close to *D. incarnata*, it can be distinguished generally by a differentially intensive maculation on leaves, and by larger, intensively coloured flowers with a more continuous and wider painting, and with a labellum with three shallow lobes. If the hybrid is more close to *D. majalis*, it can be distinguished by longer, more slender, and more erect leaves, which are terminated by hooded tips, whereby the tips of upper leaves are deflected to the stalk. The flowers are smaller, with a narrow painting.

MOLNÁR – SULYOK (1997) report about one probably extinct and one recent localities. At the Regéc locality, we found one individual.

The taxa of the genus *Dactylorhiza* grow in plant communities of the class *Molinio-Arrhenatheretea*, as documented by the following phytosociological record:

Zemplén Mts., Regéc: Gyertyánkúti-rétek, 5. 6. 1996 Hrivnák, Vlčko, Balázs, area 4 × 4 m, E<sub>1</sub> 90 %, E<sub>0</sub> 8 % (indeterminate), slope angle 3°, exposition SSW, altitude 650 m a. s. l., average height of herbs 40–48 cm.

*Festuca pratensis* 3, *Brachypodium pinnatum* 2b, *Cruciata glabra* 2a, *Dactylis glomerata* 2a, *Galium boreale* 2a, *Achillea ptarmica* 1, *Agrostis capillaris* 1, *Anthoxanthum odoratum* 1, *Betonica officinalis* 1, *Cirsium pannonicum* 1, *Filipendula vulgaris* 1, *Potentilla erecta* 1, *Prunella vulgaris* 1, *Sanguisorba officinalis* 1, *Succisa pratensis* 1, *Acetosa pratensis* +, *Achillea millefolium* +, *Avenula pubescens* +, *Briza media* +, *Campanula patula* +, *Carex montana* +, *C. pallescens* +, *C. panicea* +, *C. tomentosa* +, *Centaurea* sp. +, *Cerastium fontanum* agg. +, *Colchicum autumnale* +, *Daucus carota* +, *Deschampsia caespitosa* +, *Festuca ovina* +, *Galium verum* +, *Gladiolus imbricatus* +, *Lathyrus pratensis* +, *Luzula multiflora* +, *Lychnis flos-cuculi* +, *Myosotis scorpioides* agg. +, *Potentilla alba* +, *Primula veris* +, *Ranunculus acris* +, *R. auricomus* agg. +, *R. polyanthemus* +, *Stellaria graminea* +, *Trifolium montanum* +, *Veronica chamaedrys* +, *Viola* cf. *canina* +, *Acer campestre* r, *Ajuga reptans* r, *Dactylorhiza incarnata* subsp. *haematodes* r, *D. majalis* subsp. *majalis* r, *Juncus conglomeratus* r, *Linum catharticum* r, *Ophioglossum vulgatum* r.


In the presented record, several diagnostic species of the order *Molinetalia* and lower syntaxa occur – *Achillea ptarmica*, *Betonica officinalis*, *Carex tomentosa*, *Cirsium pannonicum*, *Deschampsia caespitosa*, *Galium boreale*, *Juncus conglomeratus*, *Ophioglossum vulgatum*, *Sanguisorba officinalis*, *Succisa pratensis*. The presence of the species of mesophilous and subxerophilous meadows and pastures (like *Festuca pratensis*, *Brachypodium pinnatum*, *Cruciata glabra*, *Dactylis glomerata*, *Agrostis capillaris*, *Anthoxanthum odoratum*, *Acetosa pratensis*, *Achillea millefolium*) indicates a substantially progressed secondary succession of wet meadows and changes of water régime.

**Acknowledgements**

The authors sincerely appreciate drawing their attention to a remarkable locality and help with field work by PELLEŠ G. from the administration of the Bükk National Park (Protected Landscape Area Zemplén). Accompanying at several interesting localities in northern Hungary by MÁRTON F., JUDIK B., BRUNDA S., FRANK T. and F. BECSEY from the Bükk National Park is also heartily acknowledged. The paper was supported by the Slovak Grant Agency for Science (Grant No. 1/7011/20).


## Summary

In the Zempléni Hegység Mts. in northeastern Hungary, we found two taxa, which are new for the Hungarian flora – *Dactylorhiza incarnata* subsp. *haematodes* var. *hyphaematodes* and *D. pulchella*, as well as a rare hybrid *D. × aschersoniana*. The authors present characteristic traits of these taxa and a phytosociological record from the locality of their occurrence.



**Fig. 1.** Map of occurrence of *Dactylorhiza incarnata* subsp. *haematodes* var. *hyphaematodes* in Hungary

**1. ábra.** A *Dactylorhiza incarnata* subsp. *haematodes* var. *hyphaematodes* előfordulása



**Fig. 2.** Map of occurrence of *Dactylorhiza pulchella* in Hungary

**2. ábra.** A *Dactylorhiza pulchella* előfordulása Magyarországon

## Összefoglalás

*Dactylorhiza incarnata* subsp. *haematodes* (Rchb.) Soó  
és *Dactylorhiza pulchella* (Druce) Averyanov Magyarországon

J. VLČKO – R. HRIVNÁK – P. BALÁZS

A Zempléni-hegységben, a Regéc község határában lévő Gyertyánkúti-réteken, 1996-ban megtaláltuk a magyarországi flórára nézve új *Dactylorhiza incarnata* subsp. *haematodes* var. *hyphaematodes*-t és a *Dactylorhiza pulchella*-t. A *Dactylorhiza pulchella* 1998-ban, a Bakonyalján, Nyirád mellett is előkerült. A *D. incarnata* subsp. *haematodes* var. *hyphaematodes* hasonlít a *D. incarnata* subsp. *incarnata*-hoz, de levelei mindkét oldalukon foltosak. A *D. pulchella* a *D. incarnata* subsp. *incarnata*-tól alacsonyabb termetével (10-25 cm) tér el, csak kivételesen érheti el az 50 cm-es magasságot, kevesebb (2-3) és keskenyebb levelével, kevésvirágú virágzatával, valamint az alacsonyabb tengerszintfeletti magasságokban 1-2 héttel későbbi virágzásával különbözik. Regéc közelében a ritka *Dactylorhiza incarnata* subsp. *incarnata* mellett feljegyeztük a *Dactylorhiza × aschersoniana* (*D. incarnata × D. majalis*) hibridet is.

## References

- BORSOS O. (1960): Geobotanische monographie der Orchideen der Pannonischen und Karpatischen Flora IV. *Dactylorhiza* I. – Ann. Univ. Sci. Budapest. Rol. Eötvös Nom. Sect. Biol. 3: 93–129.
- BARKMAN J. J. – DOING H. – SEGAL S. (1964): Kritische Bemerkungen und Vorschläge zur quantitativen Vegetationsanalyse. – Acta Bot. Neerl. (Amsterdam) 13: 394–419.
- DELFORGE P. (1994): Guide des orchidées d'Europe d'Afrique du Nord et du Proche-Orient. – Delachaux et Niestlé, Lausanne 480 pp.
- ELLMAUER T. – MUCINA L. (1993): *Molinio-Arrhenatheretea*. In: MUCINA L. – GRABHERR G. – ELLMAUER T. (eds.): Die Pflanzengesellschaften Österreichs. Teil I. Anthropogene Vegetation. – Gustav Fischer Verlag, Jena, pp. 297–401.
- MARHOLD K. – HINDÁK F. (eds., 1998): Zoznam nižších a vyšších rastlín Slovenska. – Veda, Bratislava. 687 pp.
- MOLNÁR A. – SÜLYOK J. (1997): Die Hybridorchideen Ungarns. In: VLČKO J. – HRIVNÁK R. (eds.): Európske vstavačovité (*Orchidaceae*) – výskum a ochrana. SAŽP, Banská Bystrica, pp.: 51–54.
- MATUS G. (1997): Florisztikai kutatások a zempléni Gyertyánkúti-réteken. – Kiteabelia 2(2): 313–316.
- MOLNÁR A. – SÜLYOK J. – VIDÉKI R. (1995): Vadon élő orchideák. A hazai növényvilág kincsei. – Kossuth Könyvkiadó, Bp. 160 pp.
- POTÚČEK O. (1990): Kľúč na určovanie vstavačovitých Československa. – Rosalia (mimoriadne vydanie), Nitra, 154 pp.
- SIMON T. (1977a): Vegetationsuntersuchungen im Zempléner Gebirge. – Akadémiai K., Bp. 350 pp.

SIMON T. (1977b): A Zempléni-hegység északi részének védendő flóra különlegességeiről. – *Abstracta Botanica* 5: 57-63.

SOÓ R. (1973): A magyar flóra és vegetáció rendszertani-növényföldrajzi kézikönyve V. – Akadémiai Kiadó, Budapest. pp.: 724.



A. BOROS: PLANTAE HUNGARIAE EXSICCATAE.

*Orcis incarnata* L.

Comit. Pest. In arenosis humidis  
territ. "Nagy-Csengődi erdő" pr.  
pag. Csengőd.

d. 25. mai. 1920. Alt. s. m. met. ca: 100

*Dactylorhiza pulchella* (Druce) Averyanov [Csengőd, 1920.05.25. ]  
leg.: Á. Boros, det.: A. Molnár V. Herbarium of Debrecen University (DE)